P&G[™] Purifier of Water formerly PUR Purifier of Water

P&G is working with many partners in an effort to reduce the sickness and death that result from drinking contaminated water

The Need for Clean Drinking Water

- Safe drinking water is a problem for more than 1 billion people around the world
- Waterborne diseases remain a leading cause of illness and death in the developing world
- More than 2,000 children die every day due to diarrheal diseases caused by unsafe water—more deaths than from HIV/AIDS and malaria combined

What is P&G Purifier of Water?

- P&G Purifier of Water is a powdered water purification technology packaged in a 4g sachet
- It works like a dirt magnet, pulling dirt and contaminants out of unclean water
- P&G packets use many of the same ingredients used in municipal water treatment systems
- It provides a visible signal that the water is getting cleaner, even in highly turbid water, and this
 increases user confidence
- · It also provides residual chlorination so that the water will remain usable for about a day
- The portability of the light-weight packets makes them a viable option for distribution to victims of emergencies and natural disasters or to hard-to-reach rural areas
- Only simple, readily available household implements—bucket, stir stick, filter cloth—are needed to use the P&G packets

Water Purification Process

- Add the contents of 1 packet to 10 liters (2.5 gallons) of water and stir to begin the process of precipitation and coagulation
- · Stir for 5 minutes until floc forms and the water is clear
- · Let rest for 5 minutes
- Filter water through a clean, 100% cotton cloth and dispose of separated floc in latrine
- Wait 20 minutes before drinking to allow for complete disinfection
- Store in a suitable container to prevent recontamination



Source Water



Floc Formation after P&G Packet Addition



Floc Formation after Complete Stirring



Decanting the Water through a Cloth Filter



Clean Water Ready for Storage and Use

World Health Organization (WHO)/UNICEF International Network to Promote Household Water Treatment and Safe Storage (www.who.int/household_water)

- There is conclusive evidence that simple, acceptable, low-cost interventions at the household and community level are capable of dramatically improving the microbial quality of household stored water and reducing the attendant risks of diarrheal disease and death
- Combined flocculation/disinfection systems have been classified by WHO as a "protective" technology, are effective at reducing pathogens of all classes, have demonstrated health benefits, leave residual protection, and remove muddy sediment

Results of P&G Packet Testing

Laboratory and Field Testing of P&G Packets

Effective in Removing Bacteria

	Initial	Post-
Bacteria	(org/liter)	Treatment
E. coli	2.0×10^8	ND
10 common fecal bacteria	9.2 x 10 ⁹	ND
Salmonella typhi	1.6 x 10 ⁸	ND
Vibrio cholerae	1.2 x 10 ⁸	ND
Shigella sonnei	2.2 x 10 ⁸	ND
Klebsiella terrigena	2.8 x 10 ⁸	ND
Campylobacter jejuni	2.0 x 10 ⁸	ND

ND = None Detected

Effective in Removing Viruses

_	Initial Viral	Mean
	Count/ml	Log
Virus	(log 10)	Reduction
Poliovirus	7.1	>5.0
Rotavirus	7.9	>5.0

Effective in Removing Cysts

	Mean Initial	Mean Log
Cyst	(org/liter)	Reduction
Cryptosporidium parvum	1.76 x 10 ⁶	4.0
Giardia lamblia	1.84 x 10 ⁶	3.6

Reduction of Heavy Metals

	initiai	Post-Treatment
Heavy Metal	(ppb)	(ppb)
Arsenic	229	1.2*
Chromium (III)	1300	3.1*
Lead	270	<10*

^{*} Below WHO Guidelines

Removes Organics and Some Pesticides

Test Material	Initial	Post-Treatment
Humic acid (ppm)	24-30	<1
DDT (ppb)	6	0.34

WHO Guidelines for DDT = 2 ppb

Effective* in a Wide Variety of Field Samples

Country	Initial NTU	Treated NTU
Guatemala	0 – 501	0.0 - 2.6
Kenya	0.7 - 1850	0.4 - 3.2
Morocco	0 - 244	0.0 - 1.1
Philippines	0 - 550	0.0 - 1.2
Bangladesh	10 – 35	0.0 - 1.1
South Africa	<0.2 – 54	0.2 - 0.4

^{*} Effective microbiologically as well as improving clarity




Health Intervention Trials

The U.S. Centers for Disease Control and Prevention and Johns Hopkins University have conducted 5 clinical trials of the *P&G* packets and proven that it significantly reduces diarrheal illness in children and the total population.

Location/ Setting	Study Design	Diarrhea Reduction
Rural Guatemala	2,982 people 52 weeks	24-29% reduction
Rural Guatemala	3,401 people 13 weeks	40-72% reduction
Rural Kenya turbid water	6,615 people 20 weeks	17-42% reduction
Pakistan urban setting	12,090 people 39 weeks	59-64% reduction
Liberia refugee camp	2,191 people 12 weeks	87-95% reduction

Kenyan Drinking Water Samples



Document intended for use outside of the United States